

New Value for the Rydberg Constant from the Hydrogen Balmer- β Transition

Ping Zhao^(a) and W. Lichten

Yale University, New Haven, Connecticut 06511

and

Howard Layer and James Bergquist

The National Bureau of Standards, Gaithersburg, Maryland 20899

(Received 20 November 1986)

The Rydberg constant R_∞ was determined to 3 parts in 10^{10} by direct comparison of the four H,D Balmer- β transitions with a National Bureau of Standards standard laser. This is the most precise value: $R_\infty = 109\,737.315\,73(3) \text{ cm}^{-1}$; it approaches the limits of accuracy for optical measurements. The fine-structure splittings and isotope shift are in excellent agreement with theory. The result agrees with less precise experiments by Zhao *et al.* and Biraben and Julien, but disagrees with the result of Hildum *et al.*

PACS numbers: 06.20.Jr, 32.30.Jc

The Rydberg is now the most precisely measured fundamental physical constant. R_∞ is basic to all calculations of atomic and molecular energy levels. It affects other fundamental constants via the least-squares adjustment, which it is taken as exact.

The meter is now defined exactly in terms of the speed of light $c \equiv 299\,792\,458 \text{ m/sec}^1$; wavelength and frequency are formally identical. The atomic second is defined in terms of the Cs clock: 1 sec $\equiv 9\,192\,631\,770$ periods of the ^{133}Cs ground-state hfs, to a precision in the radio-frequency region of several parts in 10^{14} . In the optical region the standard of frequency/wavelength, the $^{127}\text{I}_2$ lines at 633 nm, is given to 1.6 parts in 10^{10} .² Since hydrogen spectra occur in both regions, the Rydberg constant has the potential of tying together the frequency spectrum.

Quantum electrodynamics (QED) is one of the most important and precise of our modern theories. Balmer's formula is modified by the Dirac relativistic equation and by QED, in the Lamb shift S . Recent rf measurements of S have been quoted to a few kilohertz.^{3,4} Balmer's formula (plus Dirac and QED) should be tested with lasers to the same absolute precision, which amounts to measuring R_∞ to a few parts in 10^{12} . Such a measurement probes theoretical terms untestable by rf techniques, such as the Lamb shift in the ground state of H.

The present experiment is a step in that direction. It is an improvement on that of Amin, Caldwell, and Lichten⁵ and Zhao *et al.*,⁶ which are Balmer- α analogs of the classical Lamb-Retherford measurement of the Lamb shift.

A piezoelectrically scanned etalon had $\lambda/200$ curved ($R = 284.3 \pm 0.1 \text{ cm}$) and flat mirrors at opposite ends. These mirrors were identical to those used by one of us in a previous experiment,⁷ but were recoated to 94% reflectivity at both standard (633 nm) and unknown (486 nm) wavelengths. Unlike the previous case,⁷ the

Fresnel phase-shift errors were negligible because of the much shorter wavelength of Balmer β (486 nm) compared to the He-Ne, CH_3 laser (3390 nm).⁷

The wavelength measurements were made by the method of virtual mirrors, which eliminated corrections due to optical phase shifts in the mirror coatings. The spacings were 12.348 90, 82.039 79 cm, which gave a virtual spacing of 69.690 89 cm.

A tungsten oven was heated to 2850 K to dissociate molecular hydrogen. A beam of H (or D) atoms emerged from a slit of the oven to be bombarded by electrons and excited to the metastable $2^2S_{1/2}$, $F = I + \frac{1}{2}$ state, which lived long enough to reach a secondary-electron-emission detector. A chopped, retroflected, cw dye-laser beam at $\lambda = 486 \text{ nm}$ crossed the atomic beam at right angles and diminished the signal by exciting the $2^2S_{1/2}$ state to the $4P_{1/2}$ and $4P_{3/2}$ states. A cw He-Ne laser was offset from the standard laser with a variable frequency depending on an rf oscillator. Both offset-

TABLE I. Corrections and estimated errors, Rydberg constant (parts in 10^{10} ; average of all four H and D transitions).

Effect	Correction	Error
2S hfs	470.6	...
4P hfs	0.0	0.5
2nd order Doppler	4.3	0.9
Photon recoil	-10.2	...
Refractive index	-0.2	...
Statistical	...	1.5
Energy-level calculations	...	0.2
Standard laser	...	1.6
Stark effect, light shifts, and other minor effects	...	1.0
Root mean square error		2.6
Rounded to		3

TABLE II. Calculations for the Rydberg constant.

Transition to	Measured λ $\lambda_m - 4860 \text{ \AA}$	Corr. (10^{-10} \AA)	Corrected λ $\lambda_c - 4860 \text{ \AA}$	Calculated ^a λ $\lambda_t - 4860 \text{ \AA}$	Rydberg constant $R - 109\,737 \text{ cm}^{-1}$
H, $4P_{1/2}$	2.656 0509(5)	721.0	2.655 700 3(5)	2.655 714 94	0.315 770(12)
H, $4P_{3/2}$	2.645 2298(13)	702.8	2.644 8881 (13)	2.644 900 51	0.315 720(29)
H combined					0.315 745(25)
D, $4P_{1/2}$	1.333 0601(11)	219.8	1.332 953 3(11)	1.332 964 42	0.315 692(25)
D, $4P_{3/2}$	1.322 437(8)	214.2	1.322 139 6(8)	1.322 152 92	0.315 741(14)
D, combined					0.315 717(25)
Grand average					0.315 731(16)

^aReferences 3-6, 16, with $R_\infty = 109\,737.31544 \text{ cm}^{-1}$, $1/a = 137.035\,9895$, Lamb shift for $H = 1\,057.851\,4(19) \text{ MHz}$, for $D = 1\,059.235(27) \text{ MHz}$, $m_p/m_e = 1\,836.152\,701(37)$.

laser and dye-laser outputs were mode matched into the piezoelectrically modulated plano-concave etalon for wavelength comparison. A feedback loop locked the dye laser to an external cavity by means of the technique of Hänsch and Couillaud.⁸ Another loop locked both to the measuring etalon and a further feedback loop locked the measuring etalon to the offset laser.

Thus the frequency locking chain was formed as follows: dye laser-external cavity-measuring etalon-offset laser-standard- $^{127}\text{I}_2$ peak. Data were taken by scanning of the oscillator frequency so that every link on the chain followed and the dye laser swept across the Balmer- β transition line.

The Balmer- β frequency f is $6.17 \times 10^8 \text{ MHz}$, and the experimental linewidth Δf was 23 MHz (D) and 28 MHz (H), which gave a resolution of $f/\Delta f \approx 2.5 \times 10^7$, a threefold improvement over the previous value for Bal-

mer α .^{5,6} Although two-photon measurements in positronium⁹ and $\text{H}(1s-2s)$ ¹⁰⁻¹³ may have better resolution, the present experiment had higher precision because of lower systematic errors and direct use of the primary standard of length. This standard was identical to that used by Jennings *et al.* to establish the current definition of the meter.² As a further check on the possibility of a foreign-gas shift in the iodine cell, its Hanle effect at 502 nm was measured by Z. Zhou after the technique of Marx and Rowley.¹⁴

The atomic-beam collimation ratio was 1400 for most of the data taken. Typical operating conditions were 50% quenching of the metastables, signal-to-noise ratio of 1000/1 with a 3-sec time constant, detector current of 0.14 pA, noise 0.12 fA, 6×10^7 metastable atoms/sec, and one-way laser power 30 mW/cm^2 . The transition region was shielded magnetically ($\sim 20 \text{ mG}$ or less) and electrostatically ($\sim 1 \text{ mV/cm}$ or less). No corrections were believed necessary for Zeeman, dc or ac Stark, or polarization effects.

The theoretical isotope shifts and fine-structure separations for the H,D Balmer- β lines were used to find the correct integral order of the mirrors by the method of exact fractions.¹⁵ During a scan across the line, a simultaneous record of offset frequency and signal strength was taken digitally. The observed lines were symmetric and had a width and line shape that agreed with theoretical calculations.




FIG. 1. Comparison of recent measurements of the Rydberg constant.

TABLE III. Comparison with recent measurements for R .

Reference	$R_\infty - 109\,737 \text{ cm}^{-1}$
Amin, Caldwell, and Lichten ^a	0.31544 ± 0.00011
Hildum <i>et al.</i> ^b	0.31492 ± 0.00022
Zhao <i>et al.</i> ^c	0.31569 ± 0.00007
Biraben and co-workers ^d	0.31569 ± 0.00006
Present result	0.31573 ± 0.00003

^aReference 5.

^bReference 13.

^cReference 6.

^dReference 11.

TABLE IV. Fine structure splittings and isotope shift.

	Theory ^a	This work
Fine structure $4P_{1/2} \leftrightarrow 4P_{3/2}$		
H	1371.1294(1)	1370.9(3)
D	1371.5036(1)	1371.8(3)
Isotope shift H \leftrightarrow D		
$2S_{1/2} \leftrightarrow 4P_{1/2}$	167752.791(2)	167752.4(3)
$2S_{1/2} \leftrightarrow 4P_{3/2}$	167753.165(2)	167753.3(3)
Weighted avg.	167753.040(2)	167753.0(3)

^aFrom Reference 16.

Table I lists the systematic corrections and sources of error for the Rydberg constant. Table II gives the measured wavelengths and the calculated Rydberg constant for infinite mass: $R_{\infty} = 109737.31573(3) \text{ cm}^{-1}$, with the meter defined such that c is exactly 299792458 m/sec, and the "i" peak of $^{127}\text{I}_2$ measured to have a frequency $\nu(^3\text{He}-^{20}\text{Ne}, ^{127}\text{I}_2, i) = 473612214.789(74) \text{ MHz}$ (1.6 parts in 10^{10}).²

The error is one standard deviation (1σ). By the more conservative Comité Consultatif pour la Définition du Mètre error¹ for the wavelength standard of 3.4 parts in 10^{10} , our error changes to 4 parts in 10^{10} at the 1σ level.

Figure 1 and Table III compare recent measurement of R_{∞} . Our result is in excellent agreement with our most recent and reliable Balmer- α measurement (Zhao *et al.*)⁶ and with Biraben and co-workers,¹¹ but highly significantly disagrees with the result of Hildum *et al.*¹³

Table IV compares measured fine-structure and isotope shifts with theory.¹⁶ The agreement is excellent.

A more detailed discussion is given elsewhere.⁶

(^a)Present address: Department of Physics, Harvard University, Cambridge, MA 02138.

¹Metrologia **19** 163 (1983).

²D. A. Jennings, C. R. Pollock, F. R. Petersen, R. E. Drullinger, K. M. Evenson, J. S. Wells, J. L. Hall, and H. P. Layer, Opt. Lett. **8**, 136 (1983).

³S. R. Lundeen and F. M. Pipkin, Phys. Rev. Lett. **46**, 232 (1981).

⁴Yu. L. Sokolov and V. P. Yakovlev, Zh. Eksp. Teor. Fiz. **83**, 15 (1982) [Sov. Phys. JETP **56**, 7 (1982)]; V. G. Pal'chikov, Yu. L. Sokolov, and V. P. Yakovlev, Metrologia **21**, 99 (1985).

⁵S. R. Amin, C. D. Caldwell, and W. Lichten, Phys. Rev. Lett. **47**, 1234 (1981).

⁶P. Zhao, W. Lichten, J. C. Bergquist, and H. P. Layer, Phys. Rev. A **34**, 5138 (1986) (Balmer α), and to be published (details of Balmer- β measurements). Results were given at the annual meeting of the Optical Society of America, Seattle, Washington, 19–24 October, 1986. Ping Zhao, Ph.D. thesis, Yale University, 1986 (unpublished).

⁷H. P. Layer, R. D. Deslattes, and W. G. Schweitzer, Jr., Appl. Opt. **15**, 734 (1976).

⁸T. W. Hänsch and B. Couillaud, Opt. Commun. **35**, 441 (1980).

⁹S. Chu, A. P. Mills, Jr. and J. L. Hall, Phys. Rev. Lett. **52**, 1689 (1984).

¹⁰J. R. M. Barr, J. M. Girkin, J. M. Tolchard, and A. I. Ferguson, Phys. Rev. Lett. **56**, 580 (1986).

¹¹F. Biraben and L. Julien, Opt. Commun. **53**, 319 (1985); F. Biraben, L. Julien, and J. C. Garreau, in Proceedings of the Fourteenth International Conference on Quantum Electronics, San Francisco, California 9–13 June 1986, and unpublished results.

¹²C. J. Foot, B. Couillaud, R. G. Beausoleil, and T. W. Hänsch, Phys. Rev. Lett. **54**, 1913 (1985).

¹³E. A. Hildum, U. Boesl, D. H. McIntyre, R. G. Beausoleil, and T. W. Hänsch, Phys. Rev. Lett. **56**, 576 (1986). After this article was submitted for publication, Hildum *et al.* sent us results of a cw measurement: $R = 109737.31571(7)$, removing their discrepancy with our results.

¹⁴B. R. Marx and W. R. C. Rowley, National Physical Laboratory Report No. MOM51 1981 (unpublished); W. R. C. Rowley and B. R. Marx, Metrologia **17**, 65 (1981).

¹⁵W. Lichten, J. Opt. Soc. Am. A **2**, 1869 (1985), and **3**, 909 (1986).

¹⁶G. W. Erickson, J. Phys. Chem. Ref. Data **6**, 831 (1977).